

January 5th, 2016

**Comments to EPA
from Environmental Health Scientists and Healthcare Professionals
In support of EPA's Proposal to Revoke Chlorpyrifos
Food Residue Tolerances**

Docket ID EPA-HQ-OPP-2015-0653

We, the undersigned, write to express our strong support for EPA's proposal to revoke all food tolerances - the maximum amount of pesticide residue allowed on or in food - for the organophosphate insecticide, chlorpyrifos.¹ We urge EPA to finalize this rule (Document ID EPA-HQ-OPP-2015-0653-0001), which would result in EPA's cancelling all associated food uses of chlorpyrifos. EPA's own Revised Human Health Risk Assessment on Chlorpyrifos (2014) has acknowledged unsafe drinking water contamination.

Chlorpyrifos is a powerful developmental neurotoxicant. Exposures to even very low doses of chlorpyrifos during critical windows of exquisite vulnerability during the nine months of pregnancy and in early postnatal life can cause brain damage to children that is characterized by diminished cognitive ability (lowered IQ), problems with working memory, delays in motor development and disruptions of primitive reflexes. It should be noted that working memory skills in the early elementary school years are a strong predictor of learning outcomes and academic achievement in later years (Alloway et al. 2010). These disruptions in children's brain development appear to be permanent, irreversible and lifelong.

Extensive published science from diverse populations correlates pre-natal chlorpyrifos exposure to reduced birth weights, delayed mental and motor development in preschoolers, and reduced IQ and delays in working memory in elementary school children (Rauh et al, 2006, 2011, Whyatt et al 2005). These persistent neurocognitive findings are especially troubling. In addition, in a pilot study of 6-11 year olds, chlorpyrifos concentrations in umbilical cord blood were associated with changes in brain structure measured by magnetic resonance imaging, including cortical thinning and regional specific cortical deformations (Rauh et al, 2012).

A recent study of inner city minority children reported a link between prenatal exposure and mild to moderate arm tremors measured when the children were middle-school aged, suggesting an even broader scope of effects on the nervous system from early life exposures, and potentially latent or long term neurological damage manifesting a decade later or beyond (Rauh et al 2015). Application of chlorpyrifos to agricultural fields within 1.5 km of the home during pregnancy has also been associated with an increased incidence of autism spectrum disorders in a recent study (Shelton et al., 2014).

These epidemiologic results are consistent with data from toxicological studies which found disruption in neuronal development, neurotransmitter systems and synaptic formation as well as behavioral and cognitive impairments in test animals following low-dose perinatal chlorpyrifos exposure (Slotkin 2004; Aldridge et al. 2004, 2005; Slotkin and Seidler, 2005, Levin et al 2001; Roy et al., 2004; Garcia et al., 2002).

Associations in newborns also were seen between prenatal exposures to organophosphate pesticides generally and abnormalities in primitive reflexes, suggesting an impact on the development of the central nervous system (Engel et al, 2007; Young et al, 2005) and in children with reduction in motor function (Eskenazi et al, 2007; Rauh et al 2006; Grandjean et al 2006; Handal et 2008; Harari et al, 2010, Rauh et al, 2015), decreases in working and visual memory, processing speed, verbal comprehensive, perceptual reasoning, and full scale IQ (Bouchard et al. 2011, Engel et al, 2011, Rauh et al, 2011; Handal et al, 2008) and increases in neuropsychological problems including ADHD, pervasive developmental disorder and behaviors typical of the autism spectrum (Rauh et al, 2006, Marks et al., 2010, Furlong et al., 2014). Certain subpopulations demonstrate greater susceptibility including children of farmworkers (Castorina et al, 2010; Engel et al., 2015) and those who have reduced capacity to detoxify the OPs (Engel et al, 2015).

In addition to neurological damage, scientists have found an elevated risk of respiratory symptoms consistent with asthma (exercised-induced coughing) and persistent decreased lung function in children exposed during early-life to organophosphate pesticides (Raanan, Harley et al 2015; Raanan, Balmes et al 2015). These observations are consistent with findings in animal studies that chlorpyrifos and other OP pesticides induced airway hyper-reactivity at doses below those that caused cholinesterase inhibition (Ndlovu et al 2011; Proskocil et al 2013).

In April, 2015, extensive comments were submitted to EPA by Earthjustice, Farmworker Justice, Pesticide Action Network (PAN), Natural Resources Defense Council (NRDC) and United Farm Workers (UFW) that identified serious health concerns linked to chlorpyrifos exposure for farmworkers, their families, and bystanders.² Comments submitted to EPA by Professors R. Whyatt, T. Slotkin and D. Hattis provide a detailed analysis of serious weaknesses in EPA's use of a model provided by Dow AgroSciences, which purports to characterize cholinesterase inhibition.³ Even EPA acknowledges that the observed neurodevelopmental impacts in children from prenatal exposures occur at levels below those that cause cholinesterase inhibition in the pregnant mother.⁴

In November, 2015 EPA acknowledged that it is unable to make a safety finding regarding chlorpyrifos use as required under the Federal Food, Drug, and Cosmetic Act (FFDCA).⁵ However, prior to issuing a final rule to cancel chlorpyrifos food tolerances, EPA has said that it "plans to complete its refined drinking water analysis for the entire country as well as update its analysis of the chlorpyrifos hazard to determine whether its current regulatory approach sufficiently addresses the potential for adverse impacts on infants and children."⁶ This will lead to more delays, and continued unsafe exposures.

EPA determined 15 years ago that residential uses of chlorpyrifos resulted in exposures to children that exceeded safe levels by orders of magnitude. In 2000 EPA phased out almost all homeowner uses, but failed to extend the same protections to rural children and pregnant women who are often exposed at even higher levels through agricultural uses.⁷ Despite the scientific evidence, EPA has continued to leave rural children and the children of farmworkers in harms' way because they are exposed to chlorpyrifos through drift, volatilization, and take-home exposures from farm worker family members (Coronado et al 2011; Bradman et al, 2005; Thompson et al, 2014; Calvert et al, 2008).⁸

During these last 15 years, consumers in the U.S. have also continued to be exposed to chlorpyrifos and other organophosphate pesticides through residues on produce (Bradman et al 2015; Lu et al 2006, Vogt et al 2012).^{9 10} Children experience greater exposure to organophosphate pesticides due to their increased hand-to-mouth action, and relative to adults they eat more fruits and vegetables, drink more, and breathe more.¹¹ Thus, with each year of delay in cancelling food tolerances and agricultural and other uses of chlorpyrifos, more

children are unnecessarily at elevated risk for problems in learning, social skills, motor function, respiratory health, and other developmental domains.

We strongly urge EPA to finalize this proposal to cancel all food tolerances for chlorpyrifos, and to cancel all remaining uses of chlorpyrifos as expeditiously as possible, given the toll that chlorpyrifos takes on farmworkers, their families and their children – born and unborn, on rural and suburban families who reside near agricultural land treated with chlorpyrifos and on all Americans who consume food that has been treated with chlorpyrifos.

Respectfully,



Jennifer Sass, Ph.D.
Senior Scientist, Natural Resources Defense Council
and Professorial Lecturer, George Washington
University, Washington, DC



Robin M. Whyatt, DrPH
Professor Emeritus
Department of Environmental Health Sciences
Mailman School of Public Health, Columbia
University, New York, NY

Supporters in alphabetical order below:

The views expressed in this editorial are the opinion of the scientists who are listed below and DO NOT imply an endorsement or support for these opinions by any organizations to which they are affiliated.

Laura Anderko PhD RN
Robert and Kathleen Scanlon Endowed Chair in Values Based Health Care & Professor
Fellow, Center for Social Justice
Director, Mid-Atlantic Center for Children's Health and the Environment
School of Nursing & Health Studies
Georgetown University
Washington DC

Howard F. Andrews, PhD
Associate Professor of Biostatistics
Mailman School of Public Health, Columbia University
New York, NY

Thomas A. Arcury, PhD
Professor and Vice Chair for Research, Department of Family and Community Medicine
Director, Program in Community Engagement, Wake Forest Clinical and Translational Science Institute
Director, Center for Worker Health
Wake Forest School of Medicine
Winston-Salem, NC

David Bellinger, MSc, PhD
Professor of Neurology, Harvard Medical School
Professor in the Department of Environmental Health, Harvard T.H. Chan School of Public Health
Harvard University
Boston, MA

Charles Benbrook, PhD
Benbrook Consulting Services
Enterprise, Oregon

Deborah Bennett, PhD
Associate Professor, Division of Environmental and Occupational Health
UC Davis

Ruth Berlin, LCSW-C
Executive Director, Maryland Pesticide Education Network
Annapolis, MD

Paul Brandt-Rauf, DrPH, MD, ScD
Professor Emeritus
Environmental Health Sciences, Mailman School of Public Health
Columbia University
New York, NY

Susan Buchanan, MD, MPH
Director, Great Lakes Center for Children's Environmental Health
University of Illinois at Chicago School of Public Health
Chicago, Illinois

Sheila Bushkin-Bedient, MD, MPH
Waterford, NY

Jaehyun Byun, MD
Santa Paula Medical Clinic
Ventura, CA

David O. Carpenter, M.D.
Director, Institute for Health and the Environment University at Albany
Rensselaer, NY

Lynn Carroll, Ph.D.
Senior Scientist
TEDX, The Endocrine Disruption Exchange

Larysa Dyrszka, MD

Stephanie M. Engel, PhD
Associate Professor of Epidemiology

Gillings School of Global Public Health, University of North Carolina at Chapel Hill
Chapel Hill, NC

Pam Factor-Litvak, PhD
Professor of Epidemiology at CUMC
Associate Dean for Research Resources
Mailman School of Public Health, Columbia University
New York, NY

Linda Forst, MD, MPH
Professor, School of Public Health
University of Illinois at Chicago
Chicago, Illinois

Robert M. Gould, MD
Associate Adjunct Professor
Program on Reproductive Health and the Environment
Department of Obstetrics, Gynecology and Reproductive Sciences
UCSF School of Medicine
Immediate Past President, Physicians for Social Responsibility

Joseph H. Graziano, Ph.D.
Professor of Environmental Health Sciences and Pharmacology
Interim Chair, Department of Environmental Health Sciences
Director, The Columbia University Superfund Research Program
New York, NY 10032

Alycia Halladay, PhD
Adjunct, Rutgers University
Department of Pharmacology and Toxicology
And Chief Science Officer, Autism Science Foundation

Russ Hauser MD, ScD, MPH
Frederick Lee Hisaw Professor of Reproductive Physiology
Professor of Environmental and Occupational Epidemiology
Harvard T.H. Chan School of Public Health
Professor of Obstetrics, Gynecology and Reproductive Biology
Harvard Medical School

Irva Hertz-Picciotto, PhD
The MIND Institute (Medical Investigations of Neurodevelopmental Disorders)
Director, Environmental Health Sciences Center
Professor and Vice Chair for Research, Department of Public Health Sciences
University of California Davis School of Medicine
Davis, CA

Jane Hoppin, ScD
Associate Professor

Deputy Director, Center for Human Health and the Environment
Department of Biological Sciences
NC State University
Raleigh, NC

Katie Huffling, RN, MS, CNM
Director of Programs
Alliance of Nurses for Healthy Environments
Mount Rainier, MD

Richard J Jackson MD MPH
Professor, Fielding School of Public Health
University of California Los Angeles
Los Angeles, CA

Margaret R. Karagas, PhD
Professor & Chair, Department of Epidemiology
Geisel School of Medicine at Dartmouth

Dr Matthew Keifer MD MPH
Occupational and Internal Medicine

Jonathan Kirsch, MD
Assistant Professor, Department of Medicine
University of Minnesota

Candace Kugel, FNP, CNM, MS
Clinical Specialist
Migrant Clinicians Network
Clinton, NY

Carol F. Kwiatkowski, PhD
Executive Director
TEDX, The Endocrine Disruption Exchange
Paonia, CO
Assistant Professor Adjunct, Department of Integrative Physiology
University of Colorado at Boulder, CO

Philip J. Landrigan, MD, MSc, FAAP
Dean for Global Health
Arnhold Institute for Global Health
Professor of Preventive Medicine and Pediatrics
Icahn School of Medicine at Mount Sinai
New York, NY

Bruce P. Lanphear, MD, MPH
Clinician Scientist, Child & Family Research Institute
Professor, Faculty of Health Sciences, Simon Fraser University

Vancouver, BC

Edward D. Levin, Ph.D.
Professor of Psychiatry and Behavioral Sciences
Duke University Medical Center
Durham, NC

Amy K. Liebman, MPA, MA
Director of Environmental and Occupational Health
Migrant Clinicians Network, Salisbury, MD

Chensheng (Alex) Lu, Ph.D.
Associate Professor of Environmental Exposure Biology
Department of Environmental Health
Harvard T.H. Chan School of Public Health
Boston, MA

Emily Marquez, Ph.D.
Staff Scientist,
Pesticide Action Network North America

Linda McCauley, PhD, RN
Dean and Professor, Emory University Nell Hodgson Woodruff School of Nursing
Atlanta, GA

Rob McConnell MD
Professor of Preventive Medicine
Keck School of Medicine
University of Southern California

Kristine McVea, MD, MPH

Mark Miller MD, MPH
Assistant Clinical Professor, Pediatrics and Occupational Environmental Medicine
University of California, San Francisco

Scott Morris, MD, MPH, FACOEM
Section on Underserved Occupational Populations
American College of Occupational and Environmental Medicine

Carlos O'Bryan, M.D., FAAFP
Chief of Family Medicine Department VCMC 2015-16
Las Islas Family Medical Group
Oxnard, CA

Peter Orris, MD, MPH
Professor and Chief of Service
Occupational and Environmental Medicine

University of Illinois Hospital and Health Sciences System
835 S. Wolcott Street, Rm 144, (MC684)
Chicago, IL 60612

Nancie Payne, President
Learning Disabilities Association of America
Pittsburgh, PA

Devon Payne-Sturges, DrPH
Assistant Professor
Maryland Institute for Applied Environmental Health
University of Maryland School of Public Health
Baltimore, MD

Frederica Perera, DrPH. PhD
Professor and Director
Columbia Center for Children's Environmental Health
Mailman School of Public Health, Columbia University
New York, NY

Sara A. Quandt, PhD
Professor of Epidemiology and Prevention
Division of Public Health Sciences
Wake Forest School of Medicine
Winston-Salem, NC

Virginia A. Rauh, ScD
Professor and Vice Chair, Population and Family Health
Mailman School of Public Health, Columbia University
New York, NY

Elena Rios, MD, MSPH
President and CEO
National Hispanic Medical Association
Washington, DC 20036

James R Roberts, MD, MPH
Professor of Pediatrics
Director, South Carolina Pediatric Practice Research Network
Medical University of South Carolina

Ted Schettler MD, MPH
Science Director
Science and Environmental Health Network

Theodore Slotkin, Ph.D.
Professor, Dept. of Pharmacology & Cancer Biology

Duke University Medical Center
Durham

Rosemary Sokas, MD, MOH
Professor and Chair
Department of Human Science
Georgetown University School of Nursing and Health Studies
Washington, DC

Lisa Solinas, M.D.
Medical Director
Santa Paula Hospital Clinic
Santa Paula West Medical Group & Pediatrics

Shanna H. Swan, PhD
Professor
Department of Preventive Medicine
Icahn School of Medicine at Mount Sinai
New York, NY

Claudia Thomas, MD
New York, NY

Gayle Thomas, MD
Medical Director, NC Farmworker Health Program
Assistant Professor, UNC Family Medicine
North Carolina

Ho Luong Tran, MD, MPH
President and CEO
National Council of Asian Pacific Islander Physicians

David Wallinga, MD, MPA
Senior health officer, Natural Resources Defense Council
San Francisco, CA

Harry Wang, MD
Vice-President, Physicians for Social Responsibility/Sacramento
Clinical Professor of Psychiatry
University of California Davis School of Medicine

Minako Watabe, MD
Ventura County Medical Center
Ventura, CA

Virginia M Weaver MD MPH
Associate Professor of Environmental Health Sciences and Medicine
Associate Faculty Member, Welch Center for Prevention, Epidemiology and Clinical Research

Johns Hopkins University Blomberg School of Public Health
Baltimore, MD 21205

David H Wegman, MD, MSc
Professor Emeritus
Department of Work Environment
UMass Lowell
Lowell, MA

Mary Wolff, PhD
Professor
Department of Preventive Medicine
Icahn School of Medicine at Mount Sinai
New York, NY

Tracey J. Woodruff, PhD, MPH
Professor and Director
UCSF Program on Reproductive Health and the Environment

Sandy Young, FNP
Las Islas Family Medical Group
Oxnard, CA

Ed Zuroweste MD
Assistant Professor of Medicine, Johns Hopkins School of Medicine
Chief Medical Officer, Migrant Clinicians Network
Clinton, NY

REFERENCES:

Aldridge JE, Levin ED, Seidler FJ, Slotkin TA. Developmental exposure of rats to chlorpyrifos leads to behavioral alterations in adulthood, involving serotonergic mechanisms and resembling animal models of depression. *Environ Health Perspect.* 2005 May;113(5):527-31.

Aldridge JE, Seidler FJ, Slotkin TA. Developmental exposure to chlorpyrifos elicits sex-selective alterations of serotonergic synaptic function in adulthood: critical periods and regional selectivity for effects on the serotonin transporter, receptor subtypes, and cell signaling. *Environ Health Perspect.* 2004 Feb;112(2):148-55.

Alloway TP, Elliott J, Place M. Investigating the relationship between attention and working memory in clinical and community samples. *Child Neuropsychol.* 2010;16(3):242-54.

Bouchard MF, Chevrier J, Harley KG, Kogut K, Vedar M, Calderon N, Trujillo C, Johnson C, Bradman A, Barr DB, Eskenazi B., Prenatal exposure to organophosphate pesticides and IQ in 7-year-old children. *Environ Health Perspect.* 2011 119(8): 1189-95.

Bradman A, Eskenazi B, Barr DB, Bravo R, Castorina R, Chevrier J, Kogut K, Harnly ME, McKone TE. Organophosphate urinary metabolite levels during pregnancy and after delivery in women living in an agricultural community. *Environ Health Perspect*. 2005 Dec;113(12):1802-7.

Bradman A, Quirós-Alcalá L, Castorina R, Schall RA, Camacho J, Holland NT, Barr DB, Eskenazi B. Effect of Organic Diet Intervention on Pesticide Exposures in Young Children Living in Low-Income Urban and Agricultural Communities. *Environ Health Perspect*. 2015 Oct;123(10):1086-93.

Bradman A, Whitaker D, Quirós L, Castorina R, Claus Henn B, Nishioka M, Morgan J, Barr DB, Harnly M, Brisbin JA, Sheldon LS, McKone TE, Eskenazi B. Pesticides and their metabolites in the homes and urine of farmworker children living in the Salinas Valley, CA. *J Expo Sci Environ Epidemiol*. 2007 Jul;17(4):331-49. Epub 2006 May 31. Erratum in: *J Expo Sci Environ Epidemiol*. 2009 Nov;19(7):694-5.

Calvert GM, Karnik J, Mehler L, Beckman J, Morrissey B, Sievert J, Barrett R, Lackovic M, Mabee L, Schwartz A, Mitchell Y, Moraga-McHaley S. Acute pesticide poisoning among agricultural workers in the United States, 1998-2005. *Am J Ind Med*. 2008 Dec;51(12):883-98.

Castorina R, Bradman A, Fenster L, Barr DB, Bravo R, Vedar MG, Harnly ME, McKone TE, Eisen EA, Eskenazi B. Comparison of current-use pesticide and other toxicant urinary metabolite levels among pregnant women in the CHAMACOS cohort and NHANES. *Environ Health Perspect*. 2010 Jun;118(6):856-63.

Coronado GD, Holte S, Vigoren E, Griffith WC, Barr DB, Faustman E, Thompson B. Organophosphate pesticide exposure and residential proximity to nearby fields: evidence for the drift pathway. *J Occup Environ Med*. 2011 Aug;53(8):884-91.

Engel SM, Berkowitz GS, Barr DB, Teitelbaum SL, Siskind J, Meisel SJ, Wetmur JG, Wolff MS. Prenatal organophosphate metabolite and organochlorine levels and performance on the Brazelton Neonatal Behavioral Assessment Scale in a multiethnic pregnancy cohort. *Am J Epidemiol*. 2007 Jun 15;165(12):1397-404.

Engel SM, Wetmur J, Chen J, Zhu C, Barr DB, Canfield RL, Wolff MS. Prenatal exposure to organophosphates, paraoxonase 1, and cognitive development in childhood. *Environ Health Perspect*. 2011 Aug;119(8):1182-8

Engel SM, Bradman A, Wolff MS, Rauh VA, Harley KG, Yang JH, Hoepner LA, Barr DB, Yolton K, Vedar MG, Xu Y, Hornung RW, Wetmur JG, Chen J, Holland NT, Perera FP, Whyatt RM, Lanphear BP, Eskenazi B., Prenatal Organophosphorus Pesticide Exposure and Child Neurodevelopment at 24 Months: An Analysis of Four Birth Cohorts. *Environ Health Perspect*, 2015 Sep 29. [Epub ahead of print].

Eskenazi B, Marks AR, Bradman A, Harley K, Barr DB, Johnson C, Morga N, Jewell NP. Organophosphate pesticide exposure and neurodevelopment in young Mexican-American children. *Environ Health Perspect*. 2007 May;115(5):792-8.

Furlong MA, Engel SM, Barr DB, Wolff MS. Prenatal exposure to organophosphate pesticides and reciprocal social behavior in childhood. *Environ Int*. 2014 Sep;70:125-31.

Garcia SJ, Seidler FJ, Qiao D, Slotkin TA. Chlorpyrifos targets developing glia: effects on glial fibrillary acidic protein. *Brain Res Dev Brain Res*. 2002 Feb 28;133(2):151-61.

González-Alzaga B, Lacasaña M, Aguilar-Garduño C, Rodríguez-Barranco M, Ballester F, Rebagliato M, Hernández AF. A systematic review of neurodevelopmental effects of prenatal and postnatal organophosphate pesticide exposure. *Toxicol Lett*. 2014 Oct 15;230(2):104-21.

Grandjean P, Harari R, Barr DB, Debes F. Pesticide exposure and stunting as independent predictors of neurobehavioral deficits in Ecuadorian school children. *Pediatrics*. 2006 Mar;117(3):e546-56.

Grube A, Donaldson D, Kiely T, Wu L. Pesticides industry sales and usage: 2006 and 2007 market estimates. 2011; United States Environmental Protection Agency, Office of Pesticide Programs, Biological and Economic Analysis Division, Washington, DC, EPA 733-R-11-001

Handal AJ, Harlow SD, Breilh J, Lozoff B. Occupational exposure to pesticides during pregnancy and neurobehavioral development of infants and toddlers. *Epidemiology*. 2008 Nov;19(6):851-9.

Harari R, Julvez J, Murata K, Barr D, Bellinger DC, Debes F, Grandjean P. Neurobehavioral deficits and increased blood pressure in school-age children prenatally exposed to pesticides. *Environ Health Perspect*. 2010 Jun;118(6):890-6.

Levin ED, Addy N, Nakajima A, Christopher NC, Seidler FJ, Slotkin TA. Persistent behavioral consequences of neonatal chlorpyrifos exposure in rats. *Brain Res Dev Brain Res*. 2001 Sep 23;130(1):83-9.

Lu C, Toepel K, Irish R, Fenske RA, Barr DB, Bravo R. Organic diets significantly lower children's dietary exposure to organophosphorus pesticides. *Environ Health Perspect*. 2006 Feb;114(2):260-3.

Marks AR, Harley K, Bradman A, Kogut K, Barr DB, Johnson C, Calderon N, Eskenazi B. Organophosphate pesticide exposure and attention in young Mexican-American children: the CHAMACOS study. *Environ Health Perspect*, 2010. 118(12): 1768-74.

Proskocil BJ, Bruun DA, Jacoby DB, van Rooijen N, Lein PJ, Fryer AD. Macrophage TNF- α mediates parathion-induced airway hyperreactivity in guinea pigs. *Am J Physiol Lung Cell Mol Physiol*. 2013 Apr 15;304(8):L519-29.

Raanan R, Harley KG, Balmes JR, Bradman A, Lipsett M, Eskenazi B. Early-life exposure to organophosphate pesticides and pediatric respiratory symptoms in the CHAMACOS cohort. *Environ Health Perspect*. 2015 Feb;123(2):179-85.

Raanan R, Balmes JR, Harley KG, Gunier RB, Magzamen S, Bradman A, Eskenazi B. Decreased lung function in 7-year-old children with early-life organophosphate exposure. *Thorax*. 2015 Dec 3

Rauh VA, Arunajadai S, Horton M, Perera F, Hoepner L, Barr DB, Whyatt R. Seven-year neurodevelopmental scores and prenatal exposure to chlorpyrifos, a common agricultural pesticide. *Environ Health Perspect*. 2011 Aug;119(8):1196-201.

Rauh VA, Garfinkel R, Perera FP, Andrews HF, Hoepner L, Barr DB, Whitehead R, Tang D, Whyatt RW. Impact of prenatal chlorpyrifos exposure on neurodevelopment in the first 3 years of life among inner-city children. *Pediatrics*. 2006 Dec;118(6):e1845-59.

Rauh VA, Garcia WE, Whyatt RM, Horton MK, Barr DB, Louis ED. Prenatal exposure to the organophosphate pesticide chlorpyrifos and childhood tremor. *Neurotoxicology*. 2015 Sep 15;51:80-86.

Rauh VA, Perera FP, Horton MK, Whyatt RM, Bansal R, Hao X, Liu J, Barr DB, Slotkin TA, Peterson BS. Brain anomalies in children exposed prenatally to a common organophosphate pesticide. *Proc Natl Acad Sci U S A*. 2012 May 15;109(20):7871-6.

Roy TS, Seidler FJ, Slotkin TA. Morphologic effects of subtoxic neonatal chlorpyrifos exposure in developing rat brain: regionally selective alterations in neurons and glia. *Brain Res Dev Brain Res*. 2004 Feb 20;148(2):197-206.

Shelton JF, Geraghty EM, Tancredi DJ, Delwiche LD, Schmidt RJ, Ritz B, Hansen RL, Hertz-Picciotto I. Neurodevelopmental disorders and prenatal residential proximity to agricultural pesticides: the CHARGE study. *Environ Health Perspect*. 2014 Oct;122(10):1103-9.

Slotkin TA, Seidler FJ. The alterations in CNS serotonergic mechanisms caused by neonatal chlorpyrifos exposure are permanent. *Brain Res Dev Brain Res*. 2005 Aug 8;158(1-2):115-9.

Slotkin TA. Guidelines for developmental neurotoxicity and their impact on organophosphate pesticides: a personal view from an academic perspective. *Neurotoxicology*. 2004 Jun;25(4):631-40. Review.

Thompson B, Griffith WC, Barr DB, Coronado GD, Vigoren EM, Faustman EM. Variability in the take-home pathway: farmworkers and non-farmworkers and their children. *J Expo Sci Environ Epidemiol*. 2014 Sep-Oct;24(5):522-31

US EPA. Chlorpyrifos Revised Risk Assessment and Agreement with Registrants. 2000; Washington, DC: US Environmental Protection Agency

Vogt R, Cassady D, Frost J, Bennett DH, Hertz-Picciotto I. An assessment of exposures to toxins through diet among California residents. *Environ Health* 2012;11:83. doi: 10.1186/1476-069X-11-83
<http://www.ehjournal.net/content/11/1/83>

Young JG, Eskenazi B, Gladstone EA, Bradman A, Pedersen L, Johnson C, Barr DB, Furlong CE, Holland NT., Association between in utero organophosphate pesticide exposure and abnormal reflexes in neonates. *Neurotoxicology*, 2005. 26(2): p. 199-209.

Whyatt RM, Garfinkel R, Hoepner LA, Andrews H, Holmes D, Williams MK, Reyes A, Diaz D, Perera FP, Camann DE, Barr DB.. 2005. Biomarkers in assessing residential insecticide exposures during pregnancy and effects on fetal growth. *Toxicol Appl Pharmacol* 206:246–254.

¹ EPA proposes to revoke chlorpyrifos food residue tolerances. EPA website. October 30, 2015.
<http://www.epa.gov/pesticides/epa-proposes-revoke-chlorpyrifos-food-residue-tolerances>

² See comments by Earthjustice, Farmworker Justice, Natural Resources Defense Council, Pesticide Action Network, California Rural Legal Assistance Foundation, Farm Labor Organizing Committee, Pineros y Campesinos Unidos del Noroeste, United Farm Workers. April, 2015. Docket ID: EPA-HQ-OPP-2008-0850-0848

³ Comments by RM Whyatt, T Slotkin, D Hattis. Docket ID EPA-HQ-OPP-2008-0850-0510, EPA-HQ-OPP-2008-0850-0100, EPA-HQ-OPP-2008-0850-0092, EPA-HQ-OPP-2008-0850-0089

⁴ EPA Revised Human Health Risk Assessment on Chlorpyrifos. December 2014. Docket ID EPA-HQ-OPP-2008-0850
<http://www.epa.gov/ingredients-used-pesticide-products/revised-human-health-risk-assessment-chlorpyrifos>

⁵ EPA 2015. Proposal to revoke chlorpyrifos food residue tolerances. Website updated Nov 9, 2015.

<http://www2.epa.gov/ingredients-used-pesticide-products/proposal-revoke-chlorpyrifos-food-residue-tolerances>

⁶ EPA 2015. Proposal to revoke chlorpyrifos food residue tolerances. Website updated Nov 9, 2015.

<http://www2.epa.gov/ingredients-used-pesticide-products/proposal-revoke-chlorpyrifos-food-residue-tolerances>

⁷ <http://www.epa.gov/ingredients-used-pesticide-products/revised-human-health-risk-assessment-chlorpyrifos>

⁸ EPA ignored direct pesticide drift onto people even though direct drift poisons people at alarming rates every year.

Moreover, EPA relied on only two flawed, unpublished, non-peer-reviewed industry-sponsored studies to completely discount volatilization—a significant threat for which EPA had initially proposed buffers as large as 1000 feet. These harms disproportionately fall on farmworkers and their families, often low-income and people of color.

⁹ A dietary intervention study reported a 60% to nearly complete reduction in concentrations of two OP metabolites (malathion dicarboxylic acid (MDA), and 3,5,6-trichlor-2-pyridinol (TCPy), a metabolite of chlorpyrifos) immediately after starting the organic food diet.

¹⁰ Residues and risk associated with imported produce, and other imported specialty crops ranging from herbs and spices to tea and coffee, also account for some of the highest risk servings of food and beverages in the U.S. food supply, yet because of EPA's lack of residue data, these residues and accompanying risk have not been rigorously accounted for in chlorpyrifos dietary risk assessments.

¹¹ Approximately 75% of the general U.S. population had detectable levels of TCPy in the National Health and Nutrition Examination Survey (NHANES) from 2001-2002. Results also showed children ages 6-11 years had concentrations of TCPy (geometric mean 3.48 µg/g creatinine) two times the concentrations detected in adults (geometric mean 1.49 µg/g creatinine) (DHHS 2009) Women living in an agricultural area of California (81% had a family member who was a farmworker) had significantly higher six dialkyl phosphate (DAP) concentrations than the levels for women of similar age in the NHANES population (Bradman et al 2005)